

Matrix Multiplication

For a divide-and-conquer approach, we want an $n \times n$ matrix to look like a matrix of smaller matrices

$$\begin{pmatrix} 1 & 3 & 21 & 9 \\ 4 & 12 & 36 & 35 \\ 42 & 6 & 10 & 53 \\ 2 & 32 & 12 & 60 \end{pmatrix} \begin{pmatrix} 41 & 2 & 6 & 14 \\ 3 & 11 & 7 & 18 \\ 40 & 13 & 16 & 29 \\ 7 & 19 & 63 & 72 \end{pmatrix}$$

How is a matrix of matrices like a matrix of integers?

Ring

$$(S, +, \cdot, 0, 1)$$



Axioms

1. $+$, \cdot associate
2. $+$ commutes
3. \cdot distributes over $+$
4. 0 is identity for $+$
5. 1 is identity for \cdot
6. every a in S has an additive inverse

Example Ring

Integers modulo n , \mathbb{Z}_n

$-[a]$ is

0 is


1 is

\mathbb{Z}_4

$-[1]$ is

$[1] + [3]$

Key Example



is a ring.

Inverse of $A = (a_{ij})$ is

Get weird

$\mathbb{R}_{2, n/2}$ is

Going Between M_n and $R_{2,n/2}$

A, B, C in M_n where

A', B', C' in $R_{2,n/2}$

$$\begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix} =$$

$$\begin{pmatrix} & \\ & \end{pmatrix} = \begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix}$$

reduce $n \times n$ matrix mult to $n/2 \times n/2$ matrix ops

Complexity

Let $M(n)$ be the number of scalar operations () to multiply two $n \times n$ matrices over

Divide-and-conquer doesn't buy much

$$M(n) =$$

In general, if

$$M(n) =$$

then $M(n) \leq kn^{lg m}$ for some k ,

provided $m >$

Complexity 2

If $m = 8$, as before

$$M(n) = O(n^{\lg 8}) =$$

If $m = 7$

$$M(n) = O(n^{\lg 7})$$

Need 2×2 multiplication with 7 scalar multiplications

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} = \begin{pmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \end{pmatrix}$$

e.g., $c_{22} =$

Strassen's Algorithm

$$m_1 = (a_{12} - a_{22})(b_{21} + b_{22})$$

$$m_2 = (a_{11} + a_{22})(b_{11} + b_{22})$$

$$m_3 = (a_{11} - a_{21})(b_{11} + b_{12})$$

$$m_4 = (a_{11} + a_{12})b_{22}$$

$$m_5 = a_{11}(b_{12} - b_{22})$$

$$m_6 = a_{22}(b_{21} - b_{11})$$

$$m_7 = (a_{21} + a_{22})b_{11}$$

Don't want to use commutativity of multiplication

More Algebra

$$c_{11} = m_1 + m_2 - m_4 + m_6$$

$$c_{12} = m_4 + m_5$$

$$c_{21} = m_6 + m_7$$

$$c_{22} = m_2 - m_3 + m_5 - m_7$$

$$(a_{11}+a_{22})(b_{11}+b_{22})$$

$$- (a_{11}-a_{21})(b_{11}+b_{12})$$

$$+ a_{11}(b_{12}-b_{22})$$

$$- (a_{21}+a_{22})b_{11}$$

Another Use of Matrix of Matrices

Connect matrix inversion with matrix multiplication

Inverse of matrix A , denoted A^{-1} , is the unique matrix such that $AA^{-1} = I$

I is the identity matrix

Inverse doesn't always exist

Example

$$\begin{pmatrix} 1 & 2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} -3 & 2 \\ 2 & -1 \end{pmatrix} = \begin{pmatrix} -3+4 & 2-2 \\ -6+6 & 4-3 \end{pmatrix}$$

Inversion vs. Multiplication

Assuming matrix inversion behaves reasonably, it has the same order as multiplication

Let $I(n) =$

1. $I(n)$ in $\Omega(n^2)$
2. $I(3n)$ in $O(I(n))$

Time to multiply two $n \times n$ matrices, $M(n)$, is in $O(I(n))$.

Reduce Matrix Multiply to an Inversion Problem

Suppose we have A, B to multiply, where both are $n \times n$

Construct a $3n \times 3n$ matrix D

$$\begin{pmatrix} I_n & A & 0 \\ 0 & I_n & B \\ 0 & 0 & I_n \end{pmatrix}$$

Claim that D^{-1} is

$$\begin{pmatrix} I_n & -A & AB \\ 0 & I_n & -B \\ 0 & 0 & I_n \end{pmatrix}$$

Check

$DD^{-1} =$

$$\left(\begin{array}{ccc|ccc} \text{In} & A & 0 & \text{In} & -A & AB \\ 0 & \text{In} & B & 0 & \text{In} & -B \\ 0 & 0 & \text{In} & 0 & 0 & \text{In} \end{array} \right)$$